

MATHEMATICS

IMPORTANT FORMULAE

AND CONCEPTS

for

Summative Assessment -II

Revision

CLASS – X

2016 – 17

Prepared by

M. S. KUMARSWAMY, TGT(MATHS)

M. Sc. Gold Medallist (Elect.), B. Ed.

Kendriya Vidyalaya GaCHIBOWI i

CLASS X : CHAPTER - 4 QUADRATIC EQUATIONS

IMPORTANT FORMULAS & CONCEPTS

POLYNOMIALS

An algebraic expression of the form $p(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + \dots + a_nx^n$, where $a \neq 0$, is called a polynomial in variable x of degree n .

Here, $a_0, a_1, a_2, a_3, \dots, a_n$ are real numbers and each power of x is a non-negative integer.

e.g. $3x^2 - 5x + 2$ is a polynomial of degree 2.

$3\sqrt{x} + 2$ is not a polynomial.

➤ If $p(x)$ is a polynomial in x , the highest power of x in $p(x)$ is called **the degree of the polynomial** $p(x)$. For example, $4x + 2$ is a polynomial in the variable x of degree 1, $2y^2 - 3y + 4$ is a polynomial in the variable y of degree 2,

- ❖ A polynomial of degree 0 is called a constant polynomial.
- ❖ A polynomial $p(x) = ax + b$ of degree 1 is called a linear polynomial.
- ❖ A polynomial $p(x) = ax^2 + bx + c$ of degree 2 is called a quadratic polynomial.
- ❖ A polynomial $p(x) = ax^3 + bx^2 + cx + d$ of degree 3 is called a cubic polynomial.
- ❖ A polynomial $p(x) = ax^4 + bx^3 + cx^2 + dx + e$ of degree 4 is called a bi-quadratic polynomial.

QUADRATIC EQUATION

A polynomial $p(x) = ax^2 + bx + c$ of degree 2 is called a quadratic polynomial, then $p(x) = 0$ is known as quadratic equation.

e.g. $2x^2 - 3x + 2 = 0$, $x^2 + 5x + 6 = 0$ are quadratic equations.

METHODS TO FIND THE SOLUTION OF QUADRATIC EQUATIONS

Three methods to find the solution of quadratic equation:

1. Factorisation method
2. Method of completing the square
3. Quadratic formula method

FACTORISATION METHOD

Steps to find the solution of given quadratic equation by factorisation

- Firstly, write the given quadratic equation in standard form $ax^2 + bx + c = 0$.
- Find two numbers α and β such that sum of α and β is equal to b and product of α and β is equal to ac .
- Write the middle term bx as $\alpha x + \beta x$ and factorise it by splitting the middle term and let factors are $(x + p)$ and $(x + q)$ i.e. $ax^2 + bx + c = 0 \Rightarrow (x + p)(x + q) = 0$
- Now equate each factor to zero and find the values of x .
- These values of x are the required roots/solutions of the given quadratic equation.

METHOD OF COMPLETING THE SQUARE

Steps to find the solution of given quadratic equation by Method of completing the square:

- Firstly, write the given quadratic equation in standard form $ax^2 + bx + c = 0$.
- Make coefficient of x^2 unity by dividing all by a then we get

$$x^2 + \frac{b}{a}x + \frac{c}{a} = 0$$

- Shift the constant on RHS and add square of half of the coefficient of x i.e. $\left(\frac{b}{2a}\right)^2$ on both sides.

$$x^2 + \frac{b}{a}x = -\frac{c}{a} \Rightarrow x^2 + 2\left(\frac{b}{2a}\right)x + \left(\frac{b}{2a}\right)^2 = -\frac{c}{a} + \left(\frac{b}{2a}\right)^2$$

- Write LHS as the perfect square of a binomial expression and simplify RHS.

$$\left(x + \frac{b}{2a}\right)^2 = \frac{b^2 - 4ac}{4a^2}$$

- Take square root on both sides

$$x + \frac{b}{2a} = \pm \sqrt{\frac{b^2 - 4ac}{4a^2}}$$

- Find the value of x by shifting the constant term on RHS i.e. $x = \pm \sqrt{\frac{b^2 - 4ac}{4a^2}} - \frac{b}{2a}$

QUADRATIC FORMULA METHOD

Steps to find the solution of given quadratic equation by quadratic formula method:

- Firstly, write the given quadratic equation in standard form $ax^2 + bx + c = 0$.
- Write the values of a, b and c by comparing the given equation with standard form.
- Find discriminant $D = b^2 - 4ac$. If value of D is negative, then is no real solution i.e. solution does not exist. If value of $D \geq 0$, then solution exists follow the next step.
- Put the value of a, b and D in quadratic formula $x = \frac{-b \pm \sqrt{D}}{2a}$ and get the required roots/solutions.

NATURE OF ROOTS

The roots of the quadratic equation $ax^2 + bx + c = 0$ by quadratic formula are given by

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-b \pm \sqrt{D}}{2a}$$

where $D = b^2 - 4ac$ is called discriminant. The nature of roots depends upon the value of discriminant D. There are three cases –

Case – I

When $D > 0$ i.e. $b^2 - 4ac > 0$, then the quadratic equation has two distinct roots.

$$\text{i.e. } x = \frac{-b + \sqrt{D}}{2a} \text{ and } \frac{-b - \sqrt{D}}{2a}$$

Case – II

When $D = 0$, then the quadratic equation has two equal real roots.

$$\text{i.e. } x = \frac{-b}{2a} \text{ and } \frac{-b}{2a}$$

Case – III

When $D < 0$ then there is no real roots exist.

CLASS X : CHAPTER - 5
ARITHMETIC PROGRESSION (AP)

IMPORTANT FORMULAS & CONCEPTS

SEQUENCE

An arrangement of numbers in a definite order according to some rule is called a sequence. In other words, a pattern of numbers in which succeeding terms are obtained from the preceding term by adding/subtracting a fixed number or by multiplying with/dividing by a fixed number, is called sequence or list of numbers.

e.g. 1,2,3,4,5

A sequence is said to be finite or infinite accordingly it has finite or infinite number of terms. The various numbers occurring in a sequence are called its terms.

ARITHMETIC PROGRESSION (AP).

An **arithmetic progression** is a list of numbers in which each term is obtained by adding a fixed number to the preceding term except the first term.

This fixed number is called the common difference of the AP. It can be positive, negative or zero.

Let us denote the first term of an AP by a_1 , second term by a_2 , . . . , n th term by a_n and the common difference by d . Then the AP becomes $a_1, a_2, a_3, \dots, a_n$.

So, $a_2 - a_1 = a_3 - a_2 = \dots = a_n - a_{n-1} = d$.

The general form of an arithmetic progression is given by

$$a, a + d, a + 2d, a + 3d, \dots$$

where a is the first term and d the common difference.

n th Term of an AP

Let a_1, a_2, a_3, \dots be an AP whose first term a_1 is a and the common difference is d .

Then,

the **second term** $a_2 = a + d = a + (2 - 1) d$

the **third term** $a_3 = a_2 + d = (a + d) + d = a + 2d = a + (3 - 1) d$

the **fourth term** $a_4 = a_3 + d = (a + 2d) + d = a + 3d = a + (4 - 1) d$

.....

.....

Looking at the pattern, we can say that the **n th term** $a_n = a + (n - 1) d$.

So, **the n th term a_n of the AP with first term a and common difference d is given by**

$$a_n = a + (n - 1) d.$$

a_n is also called the **general term of the AP**. If there are m terms in the AP, then a_m represents the **last term which is sometimes also denoted by l** .

n th Term from the end of an AP

Let the last term of an AP be ' l ' and the common difference of an AP is ' d ' then the n th term from the end of an AP is given by

$$l_n = l - (n - 1) d.$$

Sum of First n Terms of an AP

The sum of the first n terms of an AP is given by

$$S_n = \frac{n}{2}[2a + (n-1)d]$$

where a = first term, d = common difference and n = number of terms.

Also, it can be written as

$$S_n = \frac{n}{2}[a + a_n]$$

where a_n = n th terms

or

$$S_n = \frac{n}{2}[a + l]$$

where l = last term

This form of the result is useful when the first and the last terms of an AP are given and the common difference is not given..

$$\text{Sum of first } n \text{ positive integers is given by } S_n = \frac{n(n+1)}{2}$$

Problems based on finding a_n if S_n is given.

Find the n th term of the AP, follow the steps:

- Consider the given sum of first n terms as S_n .
- Find the value of S_1 and S_2 by substituting the value of n as 1 and 2.
- The value of S_1 is a_1 i.e. a = first term and $S_2 - S_1 = a_2$
- Find the value of $a_2 - a_1 = d$, common difference.
- By using the value of a and d , Write AP.

Problems based on finding S_n if a_n is given.

Find the sum of n term of an AP, follow the steps:

- Consider the n th term of an AP as a_n .
- Find the value of a_1 and a_2 by substituting the value of n as 1 and 2.
- The value of a_1 is a = first term.
- Find the value of $a_2 - a_1 = d$, common difference.
- By using the value of a and d , Write AP.
- By using S_n formula, simplify the expression after substituting the value of a and d .

Arithmetic Mean

If a , b and c are in AP, then ' b ' is known as arithmetic mean between ' a ' and ' c '

$$b = \frac{a+c}{2} \text{ i.e. AM between 'a' and 'c' is } \frac{a+c}{2}.$$

CLASS X : CHAPTER - 7
COORDINATE GEOMETRY

IMPORTANT FORMULAS & CONCEPTS

Points to remember

- ☞ The distance of a point from the y -axis is called its **x -coordinate**, or **abscissa**.
- ☞ The distance of a point from the x -axis is called its **y -coordinate**, or **ordinate**.
- ☞ The coordinates of a point on the x -axis are of the form $(x, 0)$.
- ☞ The coordinates of a point on the y -axis are of the form $(0, y)$.

Distance Formula

The distance between any two points $A(x_1, y_1)$ and $B(x_2, y_2)$ is given by

$$AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

or $AB = \sqrt{(\text{difference of abscissae})^2 + (\text{difference of ordinates})^2}$

Distance of a point from origin

The distance of a point $P(x, y)$ from origin O is given by $OP = \sqrt{x^2 + y^2}$

Problems based on geometrical figure

To show that a given figure is a

- ☞ Parallelogram – prove that the opposite sides are equal
- ☞ Rectangle – prove that the opposite sides are equal and the diagonals are equal.
- ☞ Parallelogram but not rectangle – prove that the opposite sides are equal and the diagonals are not equal.
- ☞ Rhombus – prove that the four sides are equal
- ☞ Square – prove that the four sides are equal and the diagonals are equal.
- ☞ Rhombus but not square – prove that the four sides are equal and the diagonals are not equal.
- ☞ Isosceles triangle – prove any two sides are equal.
- ☞ Equilateral triangle – prove that all three sides are equal.
- ☞ Right triangle – prove that sides of triangle satisfies Pythagoras theorem.

Section formula

The coordinates of the point $P(x, y)$ which divides the line segment joining the points $A(x_1, y_1)$ and $B(x_2, y_2)$, internally, in the ratio $m_1 : m_2$ are

$$\left(\frac{m_1 x_2 + m_2 x_1}{m_1 + m_2}, \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} \right)$$

This is known as the **section formula**.

Mid-point formula


The coordinates of the point $P(x, y)$ which is the midpoint of the line segment joining the points

$A(x_1, y_1)$ and $B(x_2, y_2)$, are $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$

Area of a Triangle

If $A(x_1, y_1)$, $B(x_2, y_2)$ and $C(x_3, y_3)$ are the vertices of a ΔABC , then the area of ΔABC is given by

$$\text{Area of } \Delta ABC = \frac{1}{2}[x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)]$$



Trick to remember the formula

The formula of area of a triangle can be learn with the help of following arrow diagram:

$$\Delta ABC = \frac{1}{2} \begin{vmatrix} x_1 & y_1 \\ x_2 & y_2 \\ x_3 & y_3 \\ x_1 & y_1 \end{vmatrix}$$

Find the sum of products of numbers at the ends of the lines pointing downwards and then subtract the sum of products of numbers at the ends of the line pointing upwards, multiply the difference by

$$\frac{1}{2} . \text{.e. Area of } \Delta ABC = \frac{1}{2} [(x_1y_2 + x_2y_3 + x_3y_1) - (x_1y_3 + x_3y_2 + x_2y_1)]$$




CLASS X : CHAPTER - 7
SOME APPLICATIONS TO TRIGONOMETRY

IMPORTANT FORMULAS & CONCEPTS

ANGLE OF ELEVATION


In the below figure, the line AC drawn from the eye of the student to the top of the minar is called the *line of sight*. The student is looking at the top of the minar. The angle BAC, so formed by the line of sight with the horizontal, is called the *angle of elevation* of the top of the minar from the eye of the student. Thus, the **line of sight** is the line drawn from the eye of an observer to the point in the object viewed by the observer.



The **angle of elevation** of the point viewed is the angle formed by the line of sight with the horizontal when the point being viewed is above the horizontal level, i.e., the case when we raise our head to look at the object

ANGLE OF DEPRESSION


In the below figure, the girl sitting on the balcony is *looking down* at a flower pot placed on a stair of the temple. In this case, the line of sight is *below* the horizontal level. The angle so formed by the line of sight with the horizontal is called the *angle of depression*. Thus, the **angle of depression** of a point on the object being viewed is the angle formed by the line of sight with the horizontal when the point is below the horizontal level, i.e., the case when we lower our head to look at the point being viewed



Trigonometric Ratios (T - Ratios) of an acute angle of a right triangle

In XOY-plane, let a revolving line OP starting from OX, trace out $\angle XOP = \theta$. From P (x, y) draw $PM \perp$ to OX.

In right angled triangle OMP. OM = x (Adjacent side); PM = y (opposite side); OP = r (hypotenuse).



$$\sin \theta = \frac{\text{Opposite Side}}{\text{Hypotenuse}} = \frac{y}{r}, \quad \cos \theta = \frac{\text{Adjacent Side}}{\text{Hypotenuse}} = \frac{x}{r}, \quad \tan \theta = \frac{\text{Opposite Side}}{\text{Adjacent Side}} = \frac{y}{x}$$

$$\operatorname{cosec} \theta = \frac{\text{Hypotenuse}}{\text{Opposite Side}} = \frac{r}{y}, \quad \sec \theta = \frac{\text{Hypotenuse}}{\text{Adjacent Side}} = \frac{r}{x}, \quad \cot \theta = \frac{\text{Adjacent Side}}{\text{Opposite Side}} = \frac{x}{y}$$

Reciprocal Relations

$$\operatorname{cosec} \theta = \frac{1}{\sin \theta}, \quad \sec \theta = \frac{1}{\cos \theta} \quad \text{and} \quad \cot \theta = \frac{1}{\tan \theta}$$

Quotient Relations

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \quad \text{and} \quad \cot \theta = \frac{\cos \theta}{\sin \theta}$$

Trigonometric ratios of Complementary angles.

$$\begin{aligned} \sin(90 - \theta) &= \cos \theta & \cos(90 - \theta) &= \sin \theta \\ \tan(90 - \theta) &= \cot \theta & \cot(90 - \theta) &= \tan \theta \\ \sec(90 - \theta) &= \operatorname{cosec} \theta & \operatorname{cosec}(90 - \theta) &= \sec \theta. \end{aligned}$$

Trigonometric ratios for angle of measure.

$0^\circ, 30^\circ, 45^\circ, 60^\circ$ and 90° in tabular form.

$\angle A$	0°	30°	45°	60°	90°
sinA	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
cosA	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
tanA	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	Not defined
cosecA	Not defined	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1
secA	1	$\frac{2}{\sqrt{3}}$	$\sqrt{2}$	2	Not defined
cotA	Not defined	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0


CLASS X : CHAPTER - 10
CIRCLES

IMPORTANT FORMULAS & CONCEPTS


Circle

The collection of all the points in a plane, which are at a fixed distance from a fixed point in the plane, is called a circle.

- The fixed point is called the *centre* of the circle and the fixed distance is called the *radius* of the circle. In the below figure, O is the centre and the length OP is the radius of the circle.



- The line segment joining the centre and any point on the circle is also called a *radius* of the circle.
- A circle divides the plane on which it lies into three parts. They are: (i) inside the circle, which is also called the *interior* of the circle; (ii) the *circle* and (iii) outside the circle, which is also called the *exterior* of the circle. The circle and its interior make up the *circular region*.
- The chord is the line segment having its two end points lying on the circumference of the circle.
- The chord, which passes through the centre of the circle, is called a *diameter* of the circle.
- A *diameter* is the longest chord and all diameters have the same length, which is equal to two times the radius.
- A piece of a circle between two points is called an *arc*.
- The longer one is called the *major arc* PQ and the shorter one is called the *minor arc* PQ.
- The length of the complete circle is called its *circumference*.
- The region between a chord and either of its arcs is called a *segment* of the circular region or simply a *segment* of the circle. There are two types of segments also, which are the *major segment* and the *minor segment*.
- The region between an arc and the two radii, joining the centre to the end points of the arc is called a *sector*. The minor arc corresponds to the *minor sector* and the major arc corresponds to the *major sector*.
- In the below figure, the region OPQ is the minor sector and remaining part of the circular region is the major sector. When two arcs are equal, that is, each is a semicircle, then both segments and both sectors become the same and each is known as a *semicircular region*.



Points to Remember :

- A circle is a collection of all the points in a plane, which are equidistant from a fixed point in the plane.
- Equal chords of a circle (or of congruent circles) subtend equal angles at the centre.
- If the angles subtended by two chords of a circle (or of congruent circles) at the centre (corresponding centre) are equal, the chords are equal.
- The perpendicular from the centre of a circle to a chord bisects the chord.
- The line drawn through the centre of a circle to bisect a chord is perpendicular to the chord.
- There is one and only one circle passing through three non-collinear points.
- Equal chords of a circle (or of congruent circles) are equidistant from the centre (or corresponding centres).
- Chords equidistant from the centre (or corresponding centres) of a circle (or of congruent circles) are equal.
- If two arcs of a circle are congruent, then their corresponding chords are equal and conversely, if two chords of a circle are equal, then their corresponding arcs (minor, major) are congruent.
- Congruent arcs of a circle subtend equal angles at the centre.
- The angle subtended by an arc at the centre is double the angle subtended by it at any point on the remaining part of the circle.
- Angles in the same segment of a circle are equal.\
- Angle in a semicircle is a right angle.
- If a line segment joining two points subtends equal angles at two other points lying on the same side of the line containing the line segment, the four points lie on a circle.
- The sum of either pair of opposite angles of a cyclic quadrilateral is 180° .
- If the sum of a pair of opposite angles of a quadrilateral is 180° , then the quadrilateral is cyclic.

Secant to a Circle


A secant to a circle is a line that intersects the circle at exactly two points.

Tangent to a Circle


A tangent to a circle is a line that intersects the circle at only one point.

Given two circles, there are lines that are tangents to both of them at the same time.


☞ If the circles are separate (do not intersect), there are four possible common tangents:




☞ If the two circles touch at just one point, there are three possible tangent lines that are common to both:




☞ If the two circles touch at just one point, with one inside the other, there is just one line that is a tangent to both:



☞ If the circles overlap - i.e. intersect at two points, there are two tangents that are common to both:



☞ If the circles lie one inside the other, there are no tangents that are common to both. A tangent to the inner circle would be a secant of the outer circle.



- ☞ The tangent to a circle is perpendicular to the radius through the point of contact.
 - ☞ *The lengths of tangents drawn from an external point to a circle are equal.*
 - ☞ The centre lies on the bisector of the angle between the two tangents.
 - ☞ “If a line in the plane of a circle is perpendicular to the radius at its endpoint on the circle, then the line is tangent to the circle”.
-

CLASS X : CHAPTER - 11
CONSTRUCTIONS

IMPORTANT CONCEPTS


To construct a triangle similar to a given triangle as per given scale factor.

Example 1 - Construct a triangle similar to a given triangle ABC with its sides equal to $\frac{3}{4}$ of the corresponding sides of the triangle ABC (i.e., of scale factor $\frac{3}{4}$).

Steps of Construction :

- ☞ Draw any ray BX making an acute angle with BC on the side opposite to the vertex A.
- ☞ Locate 4 (the greater of 3 and 4 in $\frac{3}{4}$) points B₁, B₂, B₃ and B₄ on BX so that BB₁ = B₁B₂ = B₂B₃ = B₃B₄.
- ☞ Join B₄C and draw a line through B₃ (the 3rd point, 3 being smaller of 3 and 4 in $\frac{3}{4}$) parallel to B₄C to intersect BC at C'.
- ☞ Draw a line through C' parallel to the line CA to intersect BA at A' (see below figure).

Then, $\Delta A'BC'$ is the required triangle.




Example 2 : Construct a triangle similar to a given triangle ABC with its sides equal to $\frac{5}{3}$ of the corresponding sides of the triangle ABC (i.e., of scale factor $\frac{5}{3}$).

Steps of Construction :

- Draw any ray BX making an acute angle with BC on the side opposite to the vertex A.
- Locate 5 points (the greater of 5 and 3 in $\frac{5}{3}$) B₁, B₂, B₃, B₄ and B₅ on BX so that BB₁ = B₁B₂ = B₂B₃ = B₃B₄ = B₄B₅.

- Join B_3 (the 3rd point, 3 being smaller of 3 and 5 in $\frac{5}{3}$) to C and draw a line through B_5 parallel to B_3C , intersecting the extended line segment BC at C' .
- Draw a line through C' parallel to CA intersecting the extended line segment BA at A' (see the below figure).
Then $A'BC'$ is the required triangle.




To construct the tangents to a circle from a point outside it.

Given : We are given a circle with centre 'O' and a point P outside it. We have to construct two tangents from P to the circle.

Steps of construction :

- ☞ Join PO and draw a perpendicular bisector of it. Let M be the midpoint of PO.
- ☞ Taking M as centre and PM or MO as radius, draw a circle. Let it intersect the given circle at the points A and B.
- ☞ Join PA and PB.

Then PA and PB are the required two tangents.




To Construct a tangent to a circle at a given point when the centre of the circle is known.

We have a circle with centre 'O' and a point P anywhere on its circumference. Then we have to construct a tangent through P.

Steps of Construction :

- ☞ Draw a circle with centre 'O' and mark a point 'P' anywhere on it. Join OP.
- ☞ Draw a perpendicular line through the point P and name it as XY, as shown in the figure.
- ☞ XY is the required tangent to the given circle passing through P.



CLASS X : CHAPTER - 12
AREAS RELATED TO CIRCLES

IMPORTANT FORMULAS & CONCEPTS

Perimeter and Area of a Circle


Perimeter/circumference of a circle = $\pi \times \text{diameter}$
= $\pi \times 2r$ (where r is the radius of the circle)
= $2\pi r$

Area of a circle = πr^2 , where $\pi = \frac{22}{7}$

Areas of Sector and Segment of a Circle

Area of the sector of angle $\theta = \frac{\theta}{360^\circ} \times \pi r^2$, where r is the radius of the circle and θ the angle of the sector in degrees

length of an arc of a sector of angle $\theta = \frac{\theta}{360^\circ} \times 2\pi r$, where r is the radius of the circle and θ the angle of the sector in degrees



Area of the segment APB = Area of the sector OAPB – Area of Δ OAB

$$= \frac{\theta}{360^\circ} \times \pi r^2 - \text{area of } \Delta \text{ OAB}$$





☞ Area of the major sector OAQB = πr^2 – Area of the minor sector OAPB

☞ Area of major segment AQB = πr^2 – Area of the minor segment APB


☞ Area of segment of a circle = Area of the corresponding sector – Area of the corresponding triangle

CLASS X : CHAPTER - 13
SURFACE AREAS AND VOLUMES

IMPORTANT FORMULAS & CONCEPTS

S. No.	Name of the solid	Figure	Lateral / Curved surface area	Total surface area	Volume	Nomenclature
1.	Cuboid		$2h(l+b)$	$2(lb+bh+hl)$	lbh	l :length b :breadth h :height
2.	Cube		$4a^2$	$6a^2$	a^3	a :side of the cube
3.	Right prism		Perimeter of base \times height	Lateral surface area+2(area of the end surface)	area of base \times height	-
4.	Regular circular Cylinder		$2\pi rh$	$2\pi r(r+h)$	$\pi r^2 h$	r :radius of the base h :height
5.	Right pyramid		$\frac{1}{2}$ (perimeter of base) \times slant height	Lateral surfaces area+area of the base	$\frac{1}{3}$ area of the base \times height	-
6.	Right circular cone		πrl	$\pi r(l+r)$	$\frac{1}{3} \pi r^2 h$	r :radius of the base h :height l :slant height
7.	Sphere		$4\pi r^2$	$4\pi r^2$	$\frac{4}{3} \pi r^3$	r :radius
8.	Hemisphere		$2\pi r^2$	$3\pi r^2$	$\frac{2}{3} \pi r^3$	r :radius

Frustum of a Cone - If a right circular cone is cut off by a plane parallel to its base, then the portion of the cone between the cutting plane and the base of the cone is called a frustum of a cone.



Slant Height of Frustum (l)	$\sqrt{h^2 + (r_1 - r_2)^2}$
Lateral Surface Area	$\pi (r_1 + r_2) l$
Total Surface Area	$\pi \{ (r_1 + r_2) l + r_1^2 + r_2^2 \}$
Volume	$\frac{\pi}{3} (r_1^2 + r_1 r_2 + r_2^2) h$
Height of cone of which the frustum is part of (h_1)	$\frac{h r_1}{(r_1 - r_2)}$



CLASS X : CHAPTER - 15 PROBABILITY

IMPORTANT FORMULAS & CONCEPTS

PROBABILITY

Experimental or empirical probability $P(E)$ of an event E is

$$P(E) = \frac{\text{Number of trials in which the event happened}}{\text{Total number of trials}}$$

The theoretical probability (also called classical probability) of an event A , written as $P(A)$, is defined as

$$P(A) = \frac{\text{Number of outcomes favourable to A}}{\text{Number of all possible outcomes of the experiment}}$$

Two or more events of an experiment, where occurrence of an event prevents occurrences of all other events, are called **Mutually Exclusive Events**.

COMPLIMENTARY EVENTS AND PROBABILITY

We denote the event 'not E ' by \bar{E} . This is called the **complement** event of event E .

$$\text{So, } P(E) + P(\bar{E}) = 1$$

i.e., $P(E) + P(\bar{E}) = 1$, which gives us $P(\bar{E}) = 1 - P(E)$.

In general, it is true that for an event E , $P(\bar{E}) = 1 - P(E)$

- ☞ The probability of an event which is impossible to occur is 0. Such an event is called an **impossible event**.
- ☞ The probability of an event which is sure (or certain) to occur is 1. Such an event is called a **sure event** or a **certain event**.
- ☞ The probability of an event E is a number $P(E)$ such that $0 \leq P(E) \leq 1$
- ☞ An event having only one outcome is called an elementary event. The sum of the probabilities of all the elementary events of an experiment is 1.

DECK OF CARDS AND PROBABILITY

A deck of playing cards consists of 52 cards which are divided into 4 suits of 13 cards each. They are black spades (♠) red hearts (♥), red diamonds (♦) and black clubs (♣).

The cards in each suit are Ace, King, Queen, Jack, 10, 9, 8, 7, 6, 5, 4, 3 and 2. Kings, Queens and Jacks are called face cards.

Example set of 52 poker playing cards

Suit	Ace	2	3	4	5	6	7	8	9	10	Jack	Queen	King
Clubs													
Diamonds													
Hearts													
Spades													

Equally likely events : Two or more events are said to be equally likely if each one of them has an equal chance of occurrence.

Mutually Exclusive events : Two or more events are mutually exclusive if the occurrence of each event prevents the every other event.

Complementary events : Consider an event has few outcomes. Event of all other outcomes in the sample survey which are not in the favourable event is called Complementary event.

Exhaustive events : All the events are exhaustive events if their union is the sample space.

Sure events : The sample space of a random experiment is called sure or certain event as any one of its elements will surely occur in any trail of the experiment.

Impossible event : An event which will occur on any account is called an impossible event.

.....